

NONSTANDARD LAGRANGIAN SUBMANIFOLDS IN $\mathbb{CP}^n$

RIVER CHIANG

ABSTRACT. We use Hamiltonian actions to construct nonstandard (as opposed to $\mathbb{RP}^n$ and T^n) Lagrangian submanifolds of $\mathbb{CP}^n$. First of all, a quotient of $\mathbb{RP}^3$ by the dihedral group D_3 is a Lagrangian submanifold of $\mathbb{CP}^3$. Secondly, $SU(n)/\mathbb{Z}_n$ are Lagrangian submanifolds of $\mathbb{CP}^{n^2-1}$.

1. INTRODUCTION

Let (M, ω) be a symplectic manifold. A submanifold $L \subset M$ is **Lagrangian** if $\dim L = \frac{1}{2} \dim M$ and ω vanishes on $T(L)$.

Using Floer homology, Paul Biran has obtained information on the topology of Lagrangian submanifolds of some closed manifolds. One of Paul Biran's theorems concerns the Lagrangian submanifolds of $\mathbb{CP}^n$ [B-1, B-2]:

Theorem 1.1. *Let L be a Lagrangian submanifold of $\mathbb{CP}^n$ such that $H_1(L; \mathbb{Z})$ is 2-torsion (that is, $2H_1(L; \mathbb{Z}) = 0$). Then*

- (1) $H^*(L; \mathbb{Z}_2) \cong H^*(\mathbb{RP}^n; \mathbb{Z}_2)$ as graded vector spaces.
- (2) If n is even, the isomorphism in (1) is an isomorphism of graded algebras.

Seidel [Se] also has a theorem with regard to the same subject:

Theorem 1.2. *Let L be a Lagrangian submanifold of $\mathbb{CP}^n$. Then*

- (1) $H^1(L; \mathbb{Z}_{2n+2}) \neq 0$.
- (2) $H^1(L; \mathbb{Z}_{2n+2}) \not\cong (\mathbb{Z}_2)^g$ for any $g \geq 2$.
- (3) If $H^1(L; \mathbb{Z}_{2n+2}) \cong \mathbb{Z}_2$, then $H^i(L; \mathbb{Z}_2) \cong \mathbb{Z}_2$ for all $i = 0, \dots, n$.

To test the limits of these topological restriction on Lagrangian submanifolds of $\mathbb{CP}^n$, we construct new examples using Hamiltonian actions. The constructions in this paper relies on the following well-known fact:

Proposition 1.3. *Let a compact Lie group G act on a symplectic manifold (M, ω) in a Hamiltonian fashion with a moment map $\Phi: M \rightarrow \mathfrak{g}^*$. Assume $\alpha \in \mathfrak{g}^*$ is a fixed point of the coadjoint action. Then for any $m \in \Phi^{-1}(\alpha)$, the orbit $G \cdot m$ is an isotropic submanifold of M .*

Proof. An orbit is smooth. We only need to show that it is isotropic. Let ξ and η be any vector fields in the Lie algebra $\mathfrak{g}$ of G and let ξ_M and η_M be their induced vector fields on M .

$$\begin{aligned}
 \omega(m)(\xi_M(m), \eta_M(m)) &= d\langle \Phi(m), \xi \rangle(\eta_M(m)) \\
 &= \langle d\Phi(\eta_M), \xi \rangle \\
 &= \langle \eta_{\mathfrak{g}^*}(\Phi(m)), \xi \rangle \\
 &= \left\langle \frac{d}{dt} \Big|_0 \exp t\eta \cdot \alpha, \xi \right\rangle \\
 &= \langle 0, \xi \rangle
 \end{aligned}$$

Therefore $\omega(m) = 0$ for all $m \in \Phi^{-1}(\alpha)$. In particular, the orbit $G \cdot m$ is a smooth isotropic submanifold of M . $\square$

In particular, any orbit of an abelian action is isotropic as well as any orbit in the zero level set of a nonabelian action.

Example 1.4. Consider the action of the torus T^n on $\mathbb{CP}^n$ given by

$$(\lambda_1, \dots, \lambda_n) \cdot [z_0, z_1, \dots, z_n] = [z_0, \lambda_1^{-1} z_1, \dots, \lambda_n^{-1} z_n].$$

This action is Hamiltonian and its moment map is

$$\Phi([z_0, z_1, \dots, z_n]) = \frac{1}{2} \left(\frac{|z_1|^2}{\|z\|^2}, \dots, \frac{|z_n|^2}{\|z\|^2} \right),$$

where $\|z\|^2 = \sum_{i=0}^n |z_i|^2$. The free orbit T^n is a Lagrangian submanifold of $\mathbb{CP}^n$.

Our first example is a Lagrangian submanifold of $\mathbb{CP}^3$. Its topology has an interesting deviation from Theorems 1.1 and 1.2.

Theorem 1.5. *There exists a Lagrangian submanifold of $\mathbb{CP}^3$, which is a quotient of $\mathbb{RP}^3$ by the dihedral group D_3 .*

We discuss the topology of this submanifold in Section 3.

We use different actions to construct a second family of examples. It provides a generalization of the standard Lagrangian submanifold $\mathbb{RP}^3$ of $\mathbb{CP}^3$.

Theorem 1.6. *$\mathrm{SU}(n)/\mathbb{Z}_n$ is a Lagrangian submanifold of $\mathbb{CP}^{n^2-1}$.*

2. FIRST EXAMPLE

View $\mathbb{C}^4$ as the space of homogeneous polynomials of degree 3 in two variables. We can define the irreducible $\mathrm{SU}(2)$ representation $\rho: \mathrm{SU}(2) \rightarrow \mathrm{GL}(\mathbb{C}^4)$ by

$$\rho(A)(p)(z) = p(zA)$$

where $p \in \mathbb{C}^4$, $A \in \mathrm{SU}(2)$ and $z = (x, y) \in \mathbb{C}^2$. The moment map for this representation as a map from $\mathbb{C}^4$ into $\mathbb{R} \times \mathbb{C} \cong \mathfrak{su}(2)^*$ is given by

$$(u_0, u_1, u_2, u_3) \mapsto \left(\frac{3}{2}|u_0|^2 + \frac{1}{2}|u_1|^2 - \frac{1}{2}|u_2|^2 - \frac{3}{2}|u_3|^2, u_0\overline{u_1} + u_1\overline{u_2} + u_2\overline{u_3} \right).$$

So the zero level set of the moment map must satisfy

$$\begin{cases} 3|u_0|^2 + |u_1|^2 = |u_2|^2 + 3|u_3|^2, \\ u_0\overline{u_1} + u_1\overline{u_2} + u_2\overline{u_3} = 0. \end{cases}$$

This $\mathrm{SU}(2)$ action commutes with the diagonal S^1 action on $\mathbb{C}^4$, and therefore descends to an action on $\mathbb{CP}^3$.

Clearly the point $[1, 0, 0, 1]$ is in the zero level set. We know that

$$(x, y) \begin{pmatrix} \alpha & \beta \\ -\overline{\beta} & \overline{\alpha} \end{pmatrix} = (\alpha x - \overline{\beta} y, \beta x + \overline{\alpha} y),$$

where $|\alpha|^2 + |\beta|^2 = 1$. To compute the stabilizer, we set $\lambda(x^3 + y^3) = (\alpha x - \overline{\beta} y)^3 + (\beta x + \overline{\alpha} y)^3$ for any possible λ in S^1 . Direct computation shows that $\lambda = \pm 1$ when $\beta = 0$ and $\lambda = \pm i$ when $\alpha = 0$. So the stabilizer is generated by

$$(2.1) \quad \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} \quad \text{and} \quad \left\{ \begin{pmatrix} \alpha & 0 \\ 0 & \overline{\alpha} \end{pmatrix}, \alpha^6 = 1 \right\}$$

Note this $\mathrm{SU}(2)$ action is not effective on $\mathbb{CP}^3$. It induces an effective $\mathrm{SO}(3)$ action and the stabilizer for $[1, 0, 0, 1]$ is the semidirect product $\mathbb{Z}_2 \ltimes \mathbb{Z}_3$, namely, the dihedral group D_3 .

Therefore, we have a Lagrangian $L = \mathrm{SO}(3)/D_3 \cong \mathbb{RP}^3/D_3$ in $\mathbb{CP}^3$.

In fact, this is the only orbit in the zero level set. According to the Guillemin-Marle-Sternberg local normal form theorem, a neighborhood of the orbit of $[1, 0, 0, 1]$ is equivariantly symplectomorphic to a neighborhood of the zero section in $Y = \mathrm{SO}(3) \times_{D_3} \mathbb{R}^3$. It is easy to check that there exists one single orbit in the zero level set in Y . Since the zero level set is connected and since the orbit is closed, it follows that the zero level set in $\mathbb{CP}^3$ consists of one orbit only.

3. TOPOLOGY FOR THE FIRST EXAMPLE

Let Γ denote the group generated by the elements in (2.1). Then the fundamental group of L is $\pi_1(L) = \Gamma$ since $L = \mathbb{RP}^3/D_3 = S^3/\Gamma$.

Let b denote $\begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$ and a denote $\begin{pmatrix} \alpha & 0 \\ 0 & \bar{\alpha} \end{pmatrix}$, $\alpha^6 = 1$. Then $b^2 = -1$ and $ba = a^5b$. The commutator subgroup $A = \{x^{-1}y^{-1}xy \mid x, y \in \pi_1(L)\}$ is $\{1, a^2, a^4\}$. By Hurewicz Theorem, the first homology group $H_1(L; \mathbb{Z})$ is $\{1, b, b^2, b^3\} \cong \mathbb{Z}_4$.

Using the Universal Coefficient Theorem and Poincaré duality, we can calculate that

$$\begin{aligned} H_0(L; \mathbb{Z}) &= \mathbb{Z}, & H^0(L; \mathbb{Z}) &= \mathbb{Z}, & H^0(L; \mathbb{Z}_2) &= \mathbb{Z}_2, \\ H_1(L; \mathbb{Z}) &= \mathbb{Z}_4, & H^1(L; \mathbb{Z}) &= 0, & H^1(L; \mathbb{Z}_2) &= \mathbb{Z}_2, \\ H_2(L; \mathbb{Z}) &= 0, & H^2(L; \mathbb{Z}) &= \mathbb{Z}_4, & H^2(L; \mathbb{Z}_2) &= \mathbb{Z}_2, \\ H_3(L; \mathbb{Z}) &= \mathbb{Z}, & H^3(L; \mathbb{Z}) &= \mathbb{Z}, & H^3(L; \mathbb{Z}_2) &= \mathbb{Z}_2. \end{aligned}$$

Also, $H^1(L; \mathbb{Z}_8) = \mathbb{Z}_4$.

The Lagrangian L we found does not satisfy the assumption of 2-torsion as in Theorem 1.1 and Theorem 1.2(3), but $H^*(L; \mathbb{Z}_2)$ is isomorphic to $H^*(\mathbb{RP}^3; \mathbb{Z}_2)$ as graded vector spaces.

4. SECOND FAMILY OF EXAMPLES¹

Consider $\mathbb{C}^{n^2}$ as the set of n by n matrices. $\mathrm{SU}(n)$ acts naturally by left multiplication. This action is Hamiltonian and its moment map is given by

$$\Phi(Z) = \frac{i}{2} ZZ^* + \frac{1}{2i} I$$

where I is the identity matrix.

This $\mathrm{SU}(n)$ action commutes with the circle action of multiplication by λI . Therefore, it descends to an action on $\mathbb{CP}^{n^2-1}$ where the center $\mathbb{Z}_n$ of $\mathrm{SU}(n)$ acts trivially. Therefore, a free orbit in the zero level set of $\mathbb{C}^{n^2}$ descends to an orbit in the zero level set of $\mathbb{CP}^{n^2-1}$ with stabilizer $\mathbb{Z}_n$. Since the dimension of $\mathrm{SU}(n)$ is $n^2 - 1$, by Proposition 1.3, $\mathrm{SU}(n)/\mathbb{Z}_n$ is a Lagrangian submanifold of $\mathbb{CP}^{n^2-1}$.

For $n = 2$, the above-mentioned orbit $\mathrm{SU}(2)/\mathbb{Z}_2$ is the standard Lagrangian $\mathbb{RP}^3$ in $\mathbb{CP}^3$. Indeed it is the projective space of the real 4 dimensional space V spanned

¹I would like to thank Michèle Audin for showing me a beautiful alternative construction of these examples in terms of real forms of Lie groups [Au].

by the following matrices:

$$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}.$$

REFERENCES

- [Au] M. Audin, Unpublished notes.
- [A-L-P] M. Audin, F. Lalonde and L. Polterovich, *Symplectic rigidity: Lagrangian submanifolds*. In Holomorphic curves in symplectic geometry. Edited by M. Audin and J. Lafontaine. Progress in Mathematics, **117**. Birkhäuser Verlag, Basel, 1994.
- [B-1] P. Biran, *Homological uniqueness of Lagrangian submanifolds*. In preparation.
- [B-2] P. Biran, *Geometry of symplectic intersections*. Proceedings of the International Congress of Mathematicians (Beijing 2002), Vol. II, 241–255.
- [Se] P. Seidel, *Graded Lagrangian submanifolds*. Bull. Soc. Math. France **128** (2000), no. 1, 103–149.